


Berggrunnskart

Bedrock map

HONNINGSVÅG 2136-4

1:50 000




NORGES
GEOLOGISKE
UNDERSØKELSE

-NGU-

2019

Geologiske kart og data på internett: www.ngu.no



GEOLOGISCHE LINJER OG SYMBOLER

GEOLOGICAL LINES AND SYMBOLS	
/	Berggrunnsgrense; sikker / usikker eller tolket fra flyfoto / gradvis overgang Lithological boundary; certain / uncertain or interpreted from aerial photographs / gradational
—	Mindre skyverforskning, utgående av skyverforskning under flak eller skjell; Minor thrusting, based on thrusting beneath flak or scarp
—	Minst en del av skyverforskningen er tolket fra flyfoto / tolket under kvalitativt lesnæring eller vann At least part of the thrusting is interpreted from aerial photographs / interpreted below Quaternary deposits or water
—	Forkastning; sikker / usikker eller tolket fra flyfoto / tolket under kvartere lesnæring eller vann Fault; certain / uncertain or interpreted from aerial photographs / interpreted below Quaternary deposits or water
—	Bedding, med planen helling angitt (25° mot SO / lodret / vannrett) Bedding, with dip of plane indicated (25° towards SW / vertical / horizontal)
—	Rett-vei-opp / lagflagen, platt peker mot yngre lag (mot NV) Younger dipping in the succession, the arrow points towards younger layers (towards NW)
—	Primær dypslag / dyptlag, med planen helling angitt (170° mot NV / mot V) Igneous layering, with dip of plane indicated (170° towards NW / towards NE)
—	Foliasjon, skiffrighet; med planen helling angitt (60° mot NV / lodret) Foliation, scistosity; with dip of plane indicated (60° towards NW / vertical)
—	Yngre skiffrighet eller klev, med planens helling angitt (50° mot NV) Younger foliation or cleavage, with dip of plane indicated (50° towards NW)
—	Foldakse, dannet samtidig med hovedfoliasjon, med stupning angitt (20° mot NO) Fold axis, formed coevally with main foliation, with plunge indicated (20° towards NE)
—	Foldakse, dannet etter hovedfoliasjon, med stupning angitt (25° mot NO) Fold axis, postfoliated, with plunge indicated (25° towards NE)
—	Strekning / mineralisering, med stupning angitt (15° mot NO) Stretching / mineralization, with plunge indicated (15° towards NE)
—	Yngre kruslinjeasjons, med stupning angitt (15° mot NO) Younger crenulation lineation, with plunge indicated (15° towards NE)
—	Fossilsleire ¹ Fossiliferous shale
—	Steinbruddspukt Aggregate quarry
—	Snittlinje, A-A' Section line, A-A'

Aldersbestemmelse

Age determination

Prøvetakningssted 1404/39Ar-datering	
Prøvetakningssted	1404/39Ar-datering
Sample location for 1404/39Ar-dating	1404/39Ar-dating
1	Pegmatittisk gabbro: 438 ± 0.7 Ma (zirkon; magmatisme); Corfu m.fl., 2006
2	Pegmatittisk gabbro: 438 ± 0.7 Ma (zirkon; magmatisme); Corfu m.fl., 2006
3	Granittisk gang: 438 ± 0.9 Ma (zirkon; magmatisme); Corfu m.fl., 2006
4	Granittisk gang: 337.3 ± 0.4 Ma (plagioklas); Lippard og Prestvik, 1997
5	Tholeiitisk gang: 337.3 ± 0.4 Ma (plagioklas); Lippard og Prestvik, 1997

Utvælg litteratur

Selected references

- Andersen, T.B. 1979: The geology of SW Magerøy, with special emphasis on the tectono-metamorphic development. Cand. real. avhandling, University of Bergen, 338 s.
- Andersen, T.B. 1984: The stratigraphy of the Magerøy Supergroup, North Norway. Norges geologiske undersøkelse, 363, 1-23.
- Anderson, M. & S. 1998: Specialized geological mapping of the Western Caledonides. Geological Society of London Special Publications, 138, 247-265.
- Corfu, F., Andersen, T.B., Roberts, D.L. & Roberts, R.J. 2006: Early Silurian basic-ultramafic and granitic plutonism in contemporaneous felsch, Magerøy, northern Norway: U-Pb ages and regional significance. Journal of the Geological Society, London, 162, 291-301.
- Corfu, F., Gerber, M., Andersen, T.B., Torsvik, T.H. & Roberts, R.J. 2011: Age and significance of Grusvann and Silurian orogenic events in the Western Caledonides. Canadian Journal of Earth Sciences, 48, 419-440.
- Curry, C.J. 1979: A regional study of the geology of the Magerøy basic igneous complex and its envelope. Unpubl. Ph.D. thesis, University of Dundee, Scotland, 244 s.
- Hynes, S.J. 1989: The geological evolution of the Porsangerfjord area, Finnmark, Northern Norway. University Ph.D. thesis, University of Wales, Cardiff, 220 s.
- Kjersrud, K. 1985: The geology of North-western Norway, Finnmark, with special reference to the tectonometamorphic evolution. Cand. real. avhandling, University of Bergen, 338 s.
- Knott, J. & T. 1997: The geological evolution of the Porsangerfjord area, Finnmark, Northern Norway. Journal of Geology, 78, 226-2280.
- Bruton, D.L. & Harper, D.A.T. 1988: Aring-Landovøy stratigraphy and faunas across the Scandinavian Caledonides. Geological Society of London Special Publications, 38, 247-265.
- Corfu, F., Andersen, T.B., Roberts, D.L. & Roberts, R.J. 2006: Early Silurian basic-ultramafic and granitic plutonism in contemporaneous felsch, Magerøy, northern Norway: U-Pb ages and regional significance. Journal of the Geological Society, London, 162, 291-301.
- Lippard, S.J. & Prestvik, T. 1997: Carboniferous dolomite dykes on Magerøy: new age determination and regional significance. Norsk Geologisk Tidsskrift, 77, 159-166.
- Lundström, K.L., Lundström, B. & Wilson, C. & Wilson, J.R. 2002: Formation of hybrid cumulates: metacrystites in Intrusion 4 of the Honningsvåg Intrusive Suite, northern Norway. Lithos, 61, 1-19.
- Nasuti, A., Roberts, D. & Gerrigan, L. 2015: Multiphase mafic dykes in the Caledonides of northern Finnmark revealed by a new high-resolution geological model. Norwegian Journal of Geology, 95, 251-263.
- Ramsay, D.M. & Sturt, B.A. 1970: The symmetamorphic emplacement of a polymict Silurian conglomerate from Magerøy, Norway. Journal of Geology, 78, 226-2280.
- Ramsay, D.M. & Sturt, B.A. 1976: The symmetamorphic emplacement of a polymict Silurian conglomerate from Magerøy, Norway. Mineralogical Magazine, 49, 301-309.
- Roberts, D. 1998: Berggrunnskart HONNINGSVÅG – Geologisk kart over Norge, M=1:250 000. Norges geologiske undersøkelse.
- Roberts, D. 2007: Palaeocurrent data from the Kalvane Complex, northern Norway: a key element in models of terrane assembly and rotation. Norsk Geologisk Tidsskrift, 87, 297-328.
- Roberts, D. & Andersen, T.B. 1998: Norskgrønnske Beskrivelse til det berggrunnsgeologiske kartbladet. M 1:250 000. Norges geologiske undersøkelse. Skrifte, 61, 1-48.
- Roberts, D., Mitchell, J.G. & Andersen, T.B. 1991: A post-Caledonian dolomite dyke from Magerøy, northern Norway. Royal Society of Edinburgh, Philosophical Transactions, 335, 231-244.
- Robins, B., Henningsen, J. & Janzen, S. 1987: The organization and internal structure of cyclic units in the Honningsvåg Intrusive Suite, Northern Norway: a model for intrusive mechanisms, double-diffusive convection and pore-magma infiltration. In Parsons, I. (ed.) Origins of Igneous Layering. Reed, Dordrecht, 287-295.
- Stedjecka, A. & Roberts, D. 1996: Finnmark Fylke, Berggrunnsgeologi M 1:500 000. Norges geologiske undersøkelse.
- Tegner, C., Robins, B. & Sørensen, H.S. 1995: Crystallization in stratified magmas in the Honningsvåg Intrusive Suite, northern Norway: a re-appraisal. Mineralogical Magazine, 60, 41-51.
- Robins, B. 1998: The mode of emplacement of the Honningsvåg Intrusive Suite, Magerøy, northern Norway. Geological Magazine, 135, 231-244.
- Robins, B., Henningsen, J. & Janzen, S. 1987: The organization and internal structure of cyclic units in the Honningsvåg Intrusive Suite, Northern Norway: a model for intrusive mechanisms, double-diffusive convection and pore-magma infiltration. In Parsons, I. (ed.) Origins of Igneous Layering. Reed, Dordrecht, 287-295.
- Referanse til kartet: Roberts, D. 2019: Berggrunnskart HONNINGSVÅG 2136-4, M 1: 50 000. Norges geologiske undersøkelse.

Geodkjet 20.12.19 av kartkomiteen (ved Ida Gurielius), Laget for fastfjellsgeologi.

