

## Referanser

- Brunke M., Gonser T. (1997).** The ecological significance of exchange processes between rivers and groundwater. *Freshwater biology*, Vol 37, Issue 1
- Enckell H., (1969).** Distribution and dispersal of Parastenocaridae (copepoda) in northern Europe. *Oikos* 20, 493–507.
- Environment Agency (2005).** Groundwater-surface water interactions in the hyporheic zone, Science Report SC030155/SR1.
- Galassi, D. M. P., Huys, R. & Reid, J. W. (2009).** Diversity, ecology and evolution of groundwater copepods. *Freshwater Biology*, Vol 54, 691–708.
- Gibert J., Stanford J.A., Dole Olivier M.J., Ward J.V. (1994).** Basic attributes of groundwater ecosystems and prospects for research. [book auth.] Danielopol D.L., Stanford J.A., (Eds) Gibert J. *Groundwater Ecology*. London : Academic Press, 1994, pp. 7-40.
- Hahn H J, (2006).** A first approach to a quantitative ecological assessment of groundwater habitats. *The GW-Fauna-Index Limnologica* 36, 2, 119–137
- Hayashi M., Rosenberry D. O. (2002).** Effects of Ground Water Exchange on the Hydrology and Ecology of Surface Water. *Groundwater*. 40, 3.
- Iannella, M., Fiasca, B., Di Lorenzo, T., Biondi, M., Di Cicco, M., Galassi D. M. P. (2020).** Spatial distribution of stygobitic crustacean harpacticoids at the boundaries of groundwater habitat types in Europe. *Sci Rep* 10, 19043.
- Korbel, K. L., Stephenson, S. & Hose, G. C. (2019).** Sediment size influences habitat selection and use by groundwater macrofauna and meiofauna. *Aquat. Sci.* 81, 39
- Kornobis, E. (2011).** Groundwater amphipods in Iceland: Population structure and phylogenetics. PhD dissertation, Faculty of Science, University of Iceland, 116 pp.
- Olsen D.A., Young R.G. (2009).** Significance of river-aquifer interactions for reach-scale thermal patterns and trout growth potential in the Motueka River New Zealand. *Hydrogeology Journal*. 17: 175-183.
- Lamoreux, J. (2004).** Stygobites are more wide-ranging than troglobites. *J. Cave Karst. Stud.* 66, 18–19
- Särkkä J, Mäkelä J, (1998).** Troglochaetus beranecki Delachaux (Polychaeta, Archiannelida) in esker groundwaters of Finland a new class of limnic animals for northern Europe. – *Hydrobiologia* 379, 1–3, 17–2.
- Schmidt, S. I., Hahn, H.J. (2012).** What is groundwater and what does this mean to fauna? – An opinion. *Limnologica*, Vol 42., Issue 1, pp 6
- Thuli, B., Hahn, H. J. (2008).** Ecology and living conditions of groundwater fauna. Technical Report TR-08-06. Svensk Kärnbränslehantering AB. Stockholm
- United States Geological Survey, (1998).** Ground Water and Surface Water: A Single Resource. US Geological Survey Circular 1139, Denver, Colorado.