




Engerbøfjellet rutile/eclogite deposit, Naustdal, Sogn og Fjordane



TITANIUM AND IRON-TITANIUM DEPOSITS IN NORWAY

NGU FOCUS 26.02.2019 / WRITTEN BY ARE.KORNELIUSSEN@NGU.NO

Approximately 300 titanium and iron-titanium mineral occurrences and deposits are known. The most significant deposits are (1) the Tellnes ilmenite deposit (in operation) and (2) combined apatite, ilmenite and vanadium-rich titanomagnetite, both in the Rogaland Anorthosite province in southernmost Norway, and (3) the Engerbøfjellet rutile/eclogite deposit in the Sunnfjord region of western Norway.



Province/region	Type of deposit		c. age	Major deposit	% TiO ₂	Mineral resource (Mt ore)		Comments
						Identif. Reserve	Possible resource	
Arendal	Ilm-Mt (V)	igneous	1250		5-20			0.7-1.0 % V in Mt
-Bamble	Rutile	metasomatic	1180	Ødegården	2-4		> 50	> 30 ppm U in rutile
Rogaland	Ilm	igneous	925	Tellnes	18	180	> 100	
Anorthosite	Ilm	igneous	930	Storgangen	17		60	
Province	Ap, Ilm, V-Mt	igneous	930	Bjerkreim-Sokndal	4-10		> 300	c. 30 % Ap+Ilm+Mt (av. 0.7 % V)
Bergen	Ilm-Mt	igneous	950					
	Rutile	metamorphic	400	Husebø	3-6		> 50	
Sunnfjord	Ilm-Mt	igneous	1200		5-10			< 1.5 % MgO in ilm.
	Rutile	metamorphic	400	Engebøfjellet	3-4		31	> 123 < 1 ppm U in rutile
Møre	Mt(V), ilm	igneous	1700	Raudsand	4		11	> 100 25-30 % Mt (0.5 % V)
Lofoten-Vesterålen	Mt(V), ilm	igneous	1900	Selvåg	4		44	c. 30 % Mt (0.4 % V)
Other	P, Mt, Ilm	igneous	270	Kodal	3-7		15	34

1 Rutile-bearing scapolitized metagabbro.

2 Mine in operation since 1960, operated by Titania AS (Kronos Worldwide Inc.). The yearly mine production is about 500.000 t ilmenite concentrate with 44-45 % TiO₂.

3 In the Bjerkreim-Sokndal layered intrusion horizons in the magmatic sequence are distinctly enriched in apatite, ilmenite and titanomagnetite. Approximately 1 km² are regarded as potential ore with 30-35 % apatite + ilmenite + titanomagnetite (0.9 % V₂O₃).

4 The Husebø rutile-bearing eclogite deposit consists of variably eclogitized jotunite affected by retrograde alteration, with 3-5 % TiO₂ outcropping more than 100.000 m².

5 Two types of Ti deposit occur in the Sunnfjord region: magmatic ilmenite-titanomagnetite deposits associated with Palaeoproterozoic mafic intrusions, and rutile-bearing Caledonian eclogitic rocks. The ilmenite deposits in the region are of minor economic interest, while the rutile-bearing eclogites represent a major mineral resource, particularly the Engebøfjellet deposit.

6 The Rødsand deposit (in operation from 1899 to 1981) is disseminated to semi-massive Fe-Ti oxides within amphibolite. The crude ore produced contained 25-30 % magnetite, 3.5-4 % ilmenite and 0.15-0.20 % V. The magnetite concentrate contained 64% Fe, 2 % TiO₂ and 0.7 % V₂O₃, and was the raw material for production of pig iron and ferrovanadium.

7 Massive and disseminated Fe-Ti deposits occur in gabbros and anorthosites. Selvåg is the only deposit considered to be of potential economic interest, with 44 Mt probable ore with 20-30 % titanomagnetite (0.6 % V₂O₃) and minor ilmenite.

8 The larvikite-lardalite complex in the southernmost parts of the Permian Oslo Rift host a number of titanomagnetite-ilmenite-apatite concentration, of which the Kodal deposit was investigated for apatite in the 1970's and 80's, and the deposit has recently been evaluated by Kodal Minerals. The mineral resource is c. 50 Mt with c. 5 % P₂O₅ (12 % apatite), 25-40 % titanomagnetite and 5-10 % ilmenite.

MORE INFORMATION, PLEASE CONTACT:



ARE KORNELIUSSEN

are.korneliusson@ngu.no • Researcher mineral resources



HENRIK SCHIELLERUP

henrik.schiellerup@ngu.no • Director Geological Resources & Environment



GEOLOGICAL
SURVEY OF
NORWAY

- NGU -

NGU - GEOLOGICAL SURVEY OF NORWAY

CONTACT ADDRESS

PO Box 6315 Torgarden
NO-7491 Trondheim
Phone: +47 73 90 40 00

VISITING ADDRESS

Leiv Eirikssons vei 39, Trondheim
E-mail: ngu@ngu.no
www.ngu.no

SOCIAL MEDIA

Facebook /Norges.geologiske.undersokelse
Twitter /nguweb
Youtube /nguweb